

Julie E. Adams  
*Editor*



# PIP Joint Fracture Dislocations

## A Clinical Casebook

 Springer

# **PIP Joint Fracture Dislocations**



Julie E. Adams

Editor

# PIP Joint Fracture Dislocations

A Clinical Casebook



Springer

*Editor*

Julie E. Adams

Mayo Clinic

Austin, MN, USA

ISBN 978-3-319-28577-1

ISBN 978-3-319-28579-5 (eBook)

DOI 10.1007/978-3-319-28579-5

Library of Congress Control Number: 2016935288

© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

This Springer imprint is published by Springer Nature

The registered company is Springer International Publishing AG Switzerland

# Preface

Proximal interphalangeal (PIP) joint injuries are common and often unrecognized or unappreciated by the patient or physician. It is important to be able to distinguish between stable, unstable, and tenuous lesions and to understand and appreciate the various options and principles of treatment.

This clinical casebook is a practical handbook that addresses PIP joint injuries and explores principles of treatment. Some of the more common clinical scenarios that may be encountered are described. The book provides a framework for the practicing hand therapist, physician, or surgeon to understand these injuries and choose between treatment options and therapy regimens in order to lead to optimal outcomes. Attention to a stepwise surgical and rehabilitation program is provided.

Chapter topics were chosen to cover the most common and useful areas of pathology and treatment that the surgeon, physician, or therapist may encounter. Chapters are framed in terms of practical principles and case examples to enhance understanding and provide useful guidance to the clinician. Each chapter is designed to highlight clinical pearls and pitfalls and to help the clinician avoid complications and improve outcomes. Expert authors were hand-selected by the editor; these experts were specifically chosen for their expertise and experience in surgery and rehabilitation and their ability to write clear and concise value-packed chapters.

It is my hope that *PIP Fracture Dislocations: A Clinical Casebook* becomes a frequently referenced guide for practicing clinicians and residents, fellows, or students who are interested in a practical appreciation of the issues associated with PIP joint injuries. To that end, I would like to thank the expert authors who have generously contributed to this volume and shared their experience and expertise and who have made my job as editor easy and fun and the fantastic work of Patrick Carr (Springer Developmental Editor) and Kristopher Spring (Springer Publishing Editor) in keeping the work on track and bringing it to fruition. I would also like to thank my husband Scott and our daughter Sarah for their patience and unwavering support.

Austin, MN, USA

Julie E. Adams, M.D.

# Contents

<b>1 Dorsal Fracture Dislocations: Biomechanics and Management Principles .....</b>	<b>1</b>
Brandon S. Smetana and Reid W. Draeger	
<b>2 Dorsal Block Splinting for Dorsal Fracture-Dislocations.....</b>	<b>19</b>
C. Liam Dwyer, Scott W. Rogers, Alex J. Ferikes, John D. Lubahn, Terri L. Wolfe, and Katie Froehlich	
<b>3 Dorsal Block Pinning.....</b>	<b>31</b>
Erica J. Gauger and Julie E. Adams	
<b>4 Proximal Interphalangeal Joint Fracture-Dislocations: Closed Reduction Internal Fixation .....</b>	<b>43</b>
Mark A. Vitale and Robert J. Strauch	
<b>5 External Fixation of PIP Joint Dorsal Fracture-Dislocations.....</b>	<b>57</b>
Stephanie Sweet and Lawrence E. Weiss	
<b>6 Use of Force Couple Devices for Treatment of PIP Fracture-Dislocations.....</b>	<b>73</b>
Maureen O'Shaughnessy and Marco Rizzo	

<b>7</b>	<b>ORIF with Screw Fixation</b> .....	85
	Chris Lincoski	
<b>8</b>	<b>Volar Plate Arthroplasty</b> .....	93
	O. Alton Barron and Daniel S. Donovan	
<b>9</b>	<b>Pilon Fracture of the Proximal Interphalangeal Joint: Use of a Combination of Fixation Techniques</b> .....	111
	Elspeth Kinnucan	
<b>10</b>	<b>Hemi-hamate Arthroplasty</b> .....	125
	Agnes Z. Dardas and Ryan P. Calfee	
<b>11</b>	<b>Condylar Replacement Arthroplasty and Other PIP Autografts</b> .....	141
	John R. Lien and T. Greg Sommerkamp	
<b>12</b>	<b>Joint Replacement Arthroplasty: Is It Ever Indicated in PIP Joint Fracture-Dislocations?</b> .....	155
	H.L. Baltzer and Steven L. Moran	
<b>13</b>	<b>PIP Fusion</b> .....	171
	Sidney M. Jacoby and Michael P. Gaspar	
<b>14</b>	<b>Pediatric PIP Joint Injuries</b> .....	187
	Felicity G. Fishman	
<b>15</b>	<b>Nonoperative Treatment of Volar PIP Joint Fracture Dislocations</b> .....	197
	Alex J. Ferikes, Scott W. Rogers, C. Liam Dwyer, John D. Lubahn, Terri L. Wolfe, and Katie Froehlich	
<b>16</b>	<b>Operative Treatment of Volar PIP Joint Fracture-Dislocations</b> .....	211
	Lawrence E. Weiss and Stephanie Sweet	
	<b>Index</b> .....	223

# Contributors

**Julie E. Adams, MD** Mayo Clinic, Austin, MN, USA

**H.L. Baltzer, MD, FRCS(C)** Division of Plastic and Reconstructive Surgery, University of Toronto, Toronto, ON, Canada

**O. Alton Barron, MD** Department of Orthopedics, CV Starr Hand Surgery Center, Mt. Sinai—Roosevelt Hospital, Mt. Sinai Icahn School of Medicine, New York, NY, USA

**Ryan P. Calfee, MD, MSc** Department of Orthopedic Surgery, Washington University School of Medicine, St. Louis, MO, USA

**Agnes Z. Dardas, BA** Department of Orthopedic Surgery, Washington University School of Medicine, St. Louis, MO, USA

**Daniel S. Donovan, MD** Department of Orthopaedic Surgery, Mt. Sinai-St. Luke's Roosevelt, New York, NY, USA

**Reid W. Draeger, MD** Department of Orthopaedics, University of North Carolina, Chapel Hill, NC, USA

**C. Liam Dwyer, MD** Department of Orthopaedic Surgery, UPMC Hamot, Erie, PA, USA

**Alex J. Ferikes, MD** Department of Orthopaedic Surgery, UPMC Hamot, Erie, PA, USA

**Felicity G. Fishman, MD** Yale Orthopaedics and Rehabilitation, New Haven, CT, USA

**Katie Froehlich, OTR/L, CHT** Hand and Upper Body Rehabilitation Center, Erie, PA, USA

**Michael P. Gaspar, MD** The Philadelphia Hand Center, P.C., Thomas Jefferson University, Department of Orthopaedic Surgery, Philadelphia, PA, USA

**Erica J. Gauger, MD** University of Minnesota Medical Center, Minneapolis, MN, USA

**Sidney M. Jacoby, MD** The Philadelphia Hand Center, P.C., Thomas Jefferson University, Department of Orthopaedic Surgery, King of Prussia, PA, USA

**Elsbeth Kinnucan, MD** Department of Orthopaedic Surgery, Kaiser Permanente, Roseville, Roseville, CA, USA

**John R. Lien, MD** Department of Orthopaedic Surgery, University of Michigan Health Systems, Ann Arbor, MI, USA

**Chris Lincoski, MD** Department of Orthopedic Hand Surgery, University Orthopedic Center, State College, PA, USA

**John D. Lubahn, MD** Department of Orthopaedic Surgery, UPMC Hamot, Erie, PA, USA

**Steven L. Moran, MD** Department of Plastic Surgery, Mayo Clinic Hospital, Rochester, MN, USA

Department of Orthopedic Surgery, Mayo Clinic Hospital, Rochester, MN, USA

**Maureen O'Shaughnessy, MD** Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN, USA

**Marco Rizzo, MD** Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN, USA

**Scott W. Rogers, MD** Department of Orthopaedic Surgery, UPMC Hamot, Erie, PA, USA

**Brandon S. Smetana, MD** Department of Orthopaedics, University of North Carolina, Chapel Hill, NC, USA

**T. Greg Sommerkamp, MD** Tri Health- Hand Surgery Specialists, Inc., Crestview Hills, KY, USA

**Robert J. Strauch, MD** Department of Orthopaedic Surgery,  
New York-Presbyterian Medical Center, Columbia University,  
New York, NY, USA

**Stephanie Sweet, MD** Philadelphia Hand Center, Thomas  
Jefferson University, King of Prussia, PA, USA

**Mark A. Vitale, MD** ONS Foundation, Greenwich, CT, USA

**Lawrence E. Weiss, MD,** Division of Hand Surgery, OAA Hand  
Center, Lehigh Valley Hospital, Allentown, PA, USA

**Terri L. Wolfe, OTR/L, CHT** Hand and Upper Body  
Rehabilitation Center, Erie, PA, USA

# Chapter 1

## Dorsal Fracture Dislocations: Biomechanics and Management Principles

Brandon S. Smetana and Reid W. Draeger

**Abstract** Dorsal proximal interphalangeal (PIP) joint fracture dislocations represent difficult injuries to manage and are frequently missed on initial evaluation. During dorsal fracture dislocations of the PIP joint, the volar plate is either disrupted or remains attached to the volar fragments of the middle phalangeal base; thus resistance to dorsal subluxation hinges on the integrity of the bony volar articular buttress of the middle phalanx. Appropriate treatment depends on recognition of dorsal instability, if present, on lateral radiographs and maintenance of reduction throughout the healing process. Anatomic reduction of the articular surface remains less important than maintenance of stability and prevention of dorsal subluxation. As such, the most commonly utilized classification scheme divides these injuries according to the amount of volar middle phalangeal base involved in the fracture dislocation in order to guide treatment. Stable injuries involve less than 30 % of the volar middle phalangeal base and are treated with nonoperative means, typically by buddy taping or with dorsal block splinting. Unstable injuries involve more than 50 % of the volar articular base and necessitate operative intervention

---

B.S. Smetana, M.D. • R.W. Draeger, M.D. (✉)  
Department of Orthopaedics, University of North Carolina,  
3102 Bioinformatics Building, CB# 7055, Chapel Hill, NC 27599, USA  
e-mail: [bsmetana@unch.unc.edu](mailto:bsmetana@unch.unc.edu); [reid\\_draeger@med.unc.edu](mailto:reid_draeger@med.unc.edu)

to prevent dorsal subluxation. Surgical strategies include open reduction and internal fixation, percutaneous pinning, external fixation, or arthroplasty. Tenuous injuries involve 30–50 % of the volar articular base and appropriate management is less clear.

**Keywords** Dorsal proximal interphalangeal joint fracture dislocation • Biomechanics • Classification • Treatment • Management principles

## Introduction

Finger fractures are common with an annual incidence of 67.9 per 100,000 persons per year with dislocations occurring with an annual incidence of 11.2 per 1000 persons per year [1]. The proximal interphalangeal (PIP) joint sits in an unprotected position and carries a long moment arm, placing it at increased risk of injury compared to surrounding structures [2]. It is additionally susceptible to injury given the high degree of articular congruity between the proximal and middle phalanges [3]. Some injuries to the PIP joint only affect ligamentous structures or are associated with small avulsion fractures of the base of the middle phalanx at the insertion of the palmar plate, while others are associated with larger fractures of the base of the middle phalanx [2]. PIP joint fractures often go unrecognized as “sprains” or “jams” leading to delayed diagnosis. Failure to recognize and treat injuries may result in subsequent stiffness, pain, swelling, angulation, and radiographic changes associated with early arthritis [2]. When dislocations occur in conjunction with fractures, they often are significantly comminuted, can be difficult to treat, and may result in a painful and stiff joint [4]. Although much is written regarding the appropriate management of closed dorsal PIP fracture dislocations, there is wide variation regarding the choice of treatment for these injuries [1, 2, 5]. This chapter reviews the biomechanics of the PIP joint in relation to PIP fracture dislocations and the treatment principles behind the approach to their management.


## **Biomechanics**

### **Anatomy**

The PIP joint is composed of a bicondylar convex proximal phalangeal head with a central groove and a biconcave middle phalangeal articular surface with a central ridge to match the proximal phalanx [6]. The dorsal-volar length of the radial condyles is larger on the index and middle fingers; however the ulnar condyles are larger on the ring and small fingers resulting in the convergence of the fingers toward the volar scaphoid tubercle during flexion [6]. The PIP joint is mostly a simple hinge joint with the majority of motion occurring in the volar/dorsal plane [1, 4], although most refer to it as a “sloppy hinge” due to small amount of rotational and angular motion [1, 7, 8]. The range of motion of the PIP joint averages from 10° of hyperextension to 110° of flexion for a total arc of motion of 100–120° about a static center of rotation [7, 9, 10]. This long arc of motion at the PIP joint contributes to approximately 85 % of finger flexion during grasp [9]. This large range of motion necessitates limited bony constraint with the middle phalangeal articular surface covering approximately 110° of the overall 210° arc of the proximal phalangeal articular surface [10]. The axis of PIP joint rotation is a single static point, which lies on lateral radiographs at an equidistant point from the dorsal, palmar, and distal articular surface of the proximal phalanx [4]. Anatomically, this is found in between the origin of the dorsal and palmar bundles of the proper lateral collateral ligament [4].

### **Stability of the PIP Joint**

Stability of the PIP joint is provided both by the bony architecture of the proximal phalanx head and middle phalangeal base articular surfaces in addition to its soft tissue envelope. Bony constraint is provided by the cup-shaped articular surface of the middle phalangeal base on the cylindrical distal articular surface of the head of



**Fig. 1.1** Soft tissue stabilizers of the PIP joint: the “box” configuration. Soft tissue constraint to the PIP joint is afforded by a boxlike configuration composed of four distinct structures surrounding the joint: dorsally the central slip, palmarly the volar plate, and both radially and ulnarly the proper and accessory collateral ligaments (From Williams IV CS. Proximal Interphalangeal Joint Fracture Dislocations: Stable and Unstable. *Hand Clinics*. 2012; 28:409–416 p.410. Originally published In Bowers WH. The anatomy of the interphalangeal joints. In: Bowers WH, editor. *The interphalangeal joints*. New York: Churchill Livingstone; 1987. p. 11; with permission)

the proximal phalanx. This bony stability is further enhanced by the central groove separating the radial and ulnar condyles of the proximal phalanx and the corresponding central ridge of the articular surface of the middle phalangeal base.

Soft tissue stability is afforded by a “box” configuration about the hinge joint (Fig. 1.1) [1, 10]. The sides of the box are formed by the radial and ulnar proper and accessory collateral ligaments, the floor by the palmar plate, and the ceiling by the central slip [1]. The palmar plate provides constraint to both hyperextension and dorsal translation of the middle phalanx on the proximal phalanx with intact bony anatomy [10]. It arises from the distal margin of the A2 pulley and inserts at the lateral volar aspect of the middle phalanx [10]. It is composed of thick checkrein portions laterally and a thinner portion centrally [6, 10].

The proper collateral ligaments originate near the center of rotation of the PIP joint at the central aspect of the head of the proximal phalanx and course distally and volarly to insert on the volar proximal aspect of the middle phalangeal base [1, 10]. They act as second-


ary stabilizers to dorsal translation in addition to providing radial and ulnar stability, especially with the PIP joint in slight flexion [1, 11].

The accessory collateral ligaments are generally less substantial upon anatomical dissection than typically represented on pictorial depictions [1]. They originate from a more distal and volar aspect of the proximal phalanx than the origin of the proper collateral ligaments and course along a more volar trajectory inserting on the volar plate near its insertion [1, 10, 11]. The accessory collateral ligaments help to provide radial and ulnar stability to the PIP joint when in extension.

### **Injury to the PIP Joint and Instability**

Disruption of any one of the soft tissue stabilizers will not typically result in dislocation, as it typically takes injury to at least two of the structures to result in dislocation [1]. With complete disruption of the collateral ligaments and volar plate seen with dislocations, reconstruction is typically not necessary as neocollateral ligaments form and provide adequate stability to the PIP joint [12]. Lutz et al. examined this anatomical ligamentous disruption in a cadaver model during pure dislocation events. In 10° of flexion, reduction of the palmar plate to its insertion occurred, as did reduction of the collateral ligaments to their site of avulsion off of their proximal phalangeal origin [11].

Fractures can commonly occur with dislocation secondary to the transmission of a longitudinally applied force vector and a large angular moment across the joint [2]. Due to variations in force transmission, three different patterns based on the joint position at the time of injury and the injury's force vector are reported (Fig. 1.2): volar lip fractures associated with dorsal fracture dislocations, dorsal lip fractures associated with volar fracture dislocations, and pilon fractures associated with longitudinal load [2]. Hyperextension at the PIP joint leads to either palmar plate disruption near its insertion or volar avulsion injuries of the palmar lip of the middle phalanx, whereas load and shear injuries lead to more profound fractures and comminution [2].



**Fig. 1.2** PIP joint fracture dislocation: patterns of injury. (a) Palmar lip fracture and dorsal subluxation can arise from either an avulsion (hyperextension)- or an impaction shear (axial load in flexion)-type mechanism. The pattern depicted in this figure represents an impaction shear injury. (b) Dorsal lip fracture with palmar subluxation can additionally arise in a similar fashion through avulsion (hyperflexion)- or impaction shear (axial load in relative hyperextension)-type mechanisms. (c) Pilon fractures as depicted in this image occur from axial load

When Eaton first classified these injuries, he made reference to a “critical corner” encompassing the palmar plate and lateral collateral ligament attachments to the volar base of the middle phalanx, which, when disrupted, led to instability [2, 13]. He later noted that when greater than 40 % of the palmar articular surface of the middle phalanx was involved that instability resulted secondary to loss of the stabilizing effects of the palmar plate and collateral ligaments [2]. When less is involved, the remaining fibers of the collateral ligament attached to the main middle phalangeal fragment provide adequate palmar stability to prevent dorsal subluxation during extension and subsequent hinging during flexion [2]. Hastings and Carroll further support the biomechanical importance of the volar lip of the middle phalangeal base in preventing dorsal subluxation. They state that when the volar lip fragment becomes of significant size, the normal semicircular shape of the middle phalangeal base is lost leaving behind a small oblique dorsal remnant. This remnant is susceptible to the forces of the central slip in allowing dorsal subluxation about the proximal phalanx [4]. The flexor digitorum superficialis inserting on the middle phalanx additionally creates a rotational moment, exacerbating this apex dorsal deformity, and increases the tendency of the middle phalanx to hinge on the volar articular remnant during flexion [4].

This dorsal subluxation of the middle phalanx necessitates disruption of the continuity of palmar restraints including the volar plate, collateral ligaments, and the palmar bony buttress [14, 15]. The volar plate is invariably disrupted in dorsal dislocations [11], and thus, the primary determinants of stability of PIP joint dorsal fracture dislocations are the degree of volar lip involvement and the subsequent amount of remaining proper collateral ligament fibers attached to the major middle phalangeal fragment [1]. Recent biomechanical evaluation of dorsal fracture dislocations demonstrated uniform stability with involvement of only 20 % of the volar lip and instability with involvement of 60–80 % involvement. 40 % articular



**Fig. 1.2** (continued) in extension with disruption of both the dorsal and volar cortical margins (From Kiefhaber TR, Stern PJ. Clinical Perspective: Fracture Dislocations of the Proximal Interphalangeal Joint. *Journal of Hand Surgery* 1998. 23(A):368–379 p.369. Originally published by Kiefhaber TR. Phalangeal dislocations/periarticular trauma. In: Peimer CA ed. *Surgery of the hand and upper extremity*. Vol 1. New York: McGraw-Hill, 2996:963; with permission)

involvement had variable amount of instability noted on their evaluation and appeared to be the threshold for stability, although they did not examine 50 % articular involvement [16].

Work by Eaton, Hastings, Carroll, Kiefhaber, and Stern has provided evidence of the importance of the volar lip size in determining stability of the fracture dislocation and provides the basis for PIP joint fracture dislocation management.

## Management Principles

Principles of management rest on creating a concentric reduction of the middle and proximal phalanges while avoiding dorsal subluxation of the middle phalanx during terminal extension and subsequent hinging of the middle phalanx during flexion [2, 4]. Poor results have been reported with persistent subluxation [2, 4, 17]. Evidence suggests that the prevention of “hinging” of the PIP articulation is of paramount importance for optimizing PIP fracture dislocation outcomes.

Anatomic reduction of the articular surface is less important for these injuries, as long as concentricity is achieved at the PIP joint. Few investigators and leading clinicians in the field are proponents of joint surface reduction as a main goal of PIP fracture dislocation treatment [2]. There has been increasing evidence that satisfactory outcomes can be obtained without obtaining anatomic articular reduction if joint subluxation is avoided and early motion is instituted [2, 14, 18–20].

Early motion has emerged as a key component of successful management of PIP fracture dislocations and parallels the early work of Salter on the positive impact of continuous passive motion on the cartilage healing process [2, 4, 5, 21]. Although stressed in most reviews, there is a paucity of high-level evidence supporting these claims [1, 2, 4, 5]. A few series claim good outcomes with slightly delayed mobilization (with as much as 3–4 weeks of immobilization) [5], yet some report the contrary with poor outcomes found with a similar period of immobilization [1]. However, with many treatment modalities permitting and demonstrating good results with early active motion, the general consensus rests on instituting motion as early as possible provided that stability is not sacrificed.